

AFRRI SR72-1  
FEBRUARY 1972

**AFRRI**  
**SCIENTIFIC**  
**REPORT**

**PERFORMANCE OF TRAINED BEAGLES  
AFTER SUPRALETHAL DOSES  
OF RADIATION**

R. L. Chaput  
R. T. Kovacic  
E. L. Barron

**ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE**  
**Defense Nuclear Agency**  
**Bethesda, Maryland**

Approved for public release; distribution unlimited


AFRRI SR72-1

RECORD SET


All aspects of investigative programs involving the use of laboratory animals sponsored by DoD components are conducted according to the principles enunciated in the "Guide for Laboratory Animal Facilities and Care", prepared by the National Academy of Sciences - National Research Council.

PERFORMANCE OF TRAINED BEAGLES AFTER  
SUPRALETHAL DOSES OF RADIATION

R. L. CHAPUT  
R. T. KOVACIC  
E. L. BARRON



R. E. GEORGE  
Commander, MSC, USN  
Chairman  
Radiation Biology Department



MYRON I. VARON  
Captain MC USN  
Director

ARMED FORCES RADIOBIOLOGY RESEARCH INSTITUTE  
Defense Nuclear Agency  
Bethesda, Maryland

## ACKNOWLEDGMENT

The authors express their sincere appreciation to J. K. Warrenfeltz, N. L. Fleming, M. E. Flynn, and W. W. Wolfe who contributed greatly to the training, irradiation, and testing of the subjects; to A. D. Reardon, W. R. Wiegel, and J. K. Widmayer who designed and constructed the training equipment; and to C. R. Curran who provided valuable advice in establishing training procedures.

## TABLE OF CONTENTS

	Page
Foreword (Nontechnical summary) . . . . .	iii
Abstract . . . . .	v
I. Introduction . . . . .	1
II. Procedures . . . . .	1
III. Results . . . . .	4
IV. Discussion . . . . .	8
V. Conclusion . . . . .	10
References . . . . .	11

## LIST OF FIGURES

Figure 1. Beagle testing box . . . . .	2
Figure 2. Beagle performance after irradiation . . . . .	6-7

## LIST OF TABLES

Table I. Beagle Training and Performance before Irradiation . . . . .	4
Table II. Beagle Performance and Clinical Symptoms during the First Hour after Irradiation . . . . .	8

## FOREWORD

(Nontechnical summary)

For a broad range of supralethal doses of ionizing radiation, the postirradiation performance of trained monkeys, miniature pigs, and rats follows a rather well-defined pattern. Within seconds to minutes after exposure, a transient period of performance decrement occurs. Functional capabilities gradually improve so that the animals are generally performing satisfactorily within 1 hour after irradiation. Later, a rapid irreversible decline in performance leads to complete incapacitation and death.

When righting reflex and clinical symptoms were used to evaluate gross performance capabilities of untrained beagle dogs after irradiation, no obvious temporary debilitation was detected. Instead, the dogs' physical ability progressively worsened and the animals became completely incapacitated shortly before death. Unlike the other animal species which had been required to perform specific tasks so that their response could be objectively evaluated, the dog's performance was only subjectively evaluated. It was possible, therefore, that trained dogs would also show a pronounced but temporary performance decrement shortly after irradiation.

Nine beagles were trained by shock avoidance conditioning to work a discrimination problem cued by visual and auditory stimuli. The animals received 5200- to 17,600-rad midline tissue doses of pulsed mixed gamma-neutron radiation, and their postirradiation performance was evaluated.

The results indicate that the response of trained dogs to supralethal doses of radiation is different from that of monkeys, miniature pigs, and rats. Sustained

periods of temporary performance decrement did not occur in dogs during the first 60 minutes after doses of 11,900 rads or less; however, random, brief periods of performance decrement were frequent. Animals that received 17,600 rads experienced a severe irreversible performance decrement within 1 hour after irradiation. At doses below 10,000 rads dogs performed at or near preirradiation levels for several days after irradiation. Later, a rapid decline in performance led to complete incapacitation and death. At doses of 10,000 to 12,000 rads most animals performed at or near preirradiation levels for only a few hours. Thereafter, a rapid, irreversible decline in performance led to complete incapacitation and death.

## ABSTRACT

Nine beagle dogs were trained by shock avoidance conditioning to work a discrimination problem cued by visual and auditory stimuli. The animals received 5200- to 17,600-rad midline tissue doses of pulsed mixed gamma-neutron radiation, and their postirradiation performance was evaluated. During the first 60 minutes after irradiation, random, brief periods of performance decrement frequently occurred. Dogs that received less than 10,000 rads performed at or near preirradiation levels several days until they became permanently incapacitated shortly before death. After doses of 10,000 rads or more most dogs performed satisfactorily for only a few minutes to hours. Their condition then worsened precipitously and the dogs died shortly thereafter.



## I. INTRODUCTION

For a broad range of supralethal doses of ionizing radiation trained monkeys, miniature pigs, and rats experience a period of transient performance decrement shortly after irradiation.<sup>2-4</sup> Functional capabilities subsequently improve so that the animals are generally performing satisfactorily within 1 hour after irradiation. Later, a rapid irreversible decline in performance occurs and is followed by incapacitation and death.

When righting reflex and clinical symptoms were used to evaluate gross performance capabilities of untrained beagles after irradiation, no temporary debilitation was detected.<sup>1,5</sup> Instead, a progressive decline in the dogs' physical ability occurred which ultimately led to complete incapacitation shortly before death.

Since the dog's performance had only been subjectively evaluated, it was possible that the animal's ability to perform a learned task would, as in the other species, fall off temporarily almost immediately after irradiation. To test this possibility, nine trained beagles were given doses of 5200 to 17,600 rads, and their postirradiation performance was evaluated.

## II. PROCEDURES

The subjects were nine male AKC registrable beagles. They were 1-1/2 to 3 years old and weighed 8 to 12 kg at the time of irradiation.

The dogs were trained to briefly flex the appropriate forelimb on cue. The animals were placed in the testing box shown in Figure 1 where only the head was firmly restrained by an adjustable yoke. The front paws were strapped to metallic grid keys which, when manipulated, actuated electrical circuits that served to record

forelimb flexing. The keys also functioned as the electrodes for the shock avoidance conditioning.

A high (1125 Hz) and a low (450 Hz) frequency tone from a 2-inch diameter speaker directly above the dog's head provided the auditory cues, and a 12-volt incandescent light in front of each key provided the visual cue. The dogs were trained to flex the right forelimb in response to the light in front of the right key and the high frequency tone and to flex the left forelimb in response to the light in front of the left key and the low frequency tone.

The cues were started at the beginning of each 8- or 10-second trial. If the dog responded correctly by flexing the appropriate forelimb within 6 seconds, it avoided electrical shock, and the cues were terminated. The remainder of the trial




Figure 1. Beagle testing box

served as a rest period. When the dog responded incorrectly by flexing the wrong forelimb, the cues were terminated, and a 1/2-second electrical shock was applied through the grid key to the paw of the forelimb that should have been flexed. If, after 6 seconds, the dog did not respond (scored as an omission) the cues were turned off and an electrical shock was delivered.

Each dog was trained until its performance reached a steady-state level. Eight dogs achieved a 90 percent or more correct response steady-state level; one dog performed consistently at the 85 percent correct response level but was retained in the study. Each animal performed at its steady-state level for approximately 2000 trials before irradiation. These were given in 100-trial sessions over 10 days.

Beagle performance before irradiation is shown in Table I. After the animals' performance levels had been established, omissions were rare.

The dogs were easily distracted by sudden changes in their surroundings. Note the slight decline in base-line performance of most animals when transferred from the training room to the exposure room. In spite of differences in the duration of training, the data in Table I indicate that the animals were equally well trained.

The dogs were exposed unilaterally to the left side to a pulse of mixed gamma-neutron radiation from the AFRRI-TRIGA reactor.<sup>5</sup> Midline tissue doses to the head were 5200 to 17,600 rads. During irradiation the dogs were held in the Plexiglas testing box (Figure 1). The animals were positioned so that the brain center line was 60 or 100 cm from the vertical reactor core center line. All doses were calculated by previously reported methods.<sup>3</sup>

Table I. Beagle Training and Performance before Irradiation

Dog #	Steady-state performance* before irradiation		Preirradiation base-line performance† in training room			Exposure room
	Total time (days)	Total number of trials	Number of trials	Percent correct response $\pm$ S.D.	Latency $\pm$ S.D. (seconds/trial)	Percent correct response
1	20	2600	2600	94.8 $\pm$ 4.0	0.93 $\pm$ 0.204	95.6
2	46	7100	2000	98.9 $\pm$ 1.1	1.37 $\pm$ 0.260	95.6
3	54	7800	2100	97.0 $\pm$ 3.0	0.58 $\pm$ 0.104	97.0
4	47	5800	2100	95.7 $\pm$ 4.13	0.380 $\pm$ 0.082	92.0
5	19	1900	1900	94.8 $\pm$ 3.1	0.607 $\pm$ 0.149	92.0
6	7	1800	1800	96.1 $\pm$ 1.5	0.682 $\pm$ 0.114	89.0
7	18	2500	2500	85.1 $\pm$ 9.90	0.126 $\pm$ 0.066	84.0
8	65	10,900	3050	97.4 $\pm$ 2.10	0.388 $\pm$ 0.098	88.6
9	19	3400	2800	94.1 $\pm$ 3.11	0.348 $\pm$ 0.121	96.0

\* 85% correct response for dog 7; 90% correct response for all other dogs

† Values based on the last trials performed before irradiation

Within 1 hour before irradiation the animals were tested in the exposure room for from one to three 100-trial sessions. After irradiation the dogs were tested in 100-trial sessions starting at 0, 20, and 40 minutes postirradiation. Thereafter, they were tested at least once a day until death. Several dogs that received higher doses were also tested at additional times during the first 6 hours after irradiation.

### III. RESULTS

Postirradiation performance data for each of the nine dogs are illustrated in Figure 2. Each bar represents the results of a test session which lasted for the times shown on the abscissa. For the two dogs that received 5200 rads, the test session immediately after the pulse consisted of a series of 70 trials. All other test sessions in this study consisted of a series of 100 trials.

For the first 60 minutes after irradiation correct responses were also averaged over 10-trial sessions. These values are shown by the data points on the solid lines.

Beagle performance and clinical symptoms for the first hour after irradiation are summarized in Table II. For purposes of this study, performance decrement was defined as "slight" for 70 or 80 percent correct response and "severe" for 60 percent correct response or less.

As shown by Figure 2, dogs that received doses of 11,900 rads or less generally performed well during the initial 60 minutes after irradiation, whereas dogs that received 17,600 rads experienced severe irreversible performance decrement within 1 hour after irradiation. This is clearly shown by the data in Table II; the dogs that received 17,600 rads had a marked increase in the number of 10-trial sessions with performance decrement compared to dogs that received 11,900 rads or less.

Although dogs 4 (7,800 rads), 5 (10,200 rads), 6 (10,200 rads), 7 (11,900 rads), and 9 (17,600 rads) showed signs of an early transient performance decrement within 15 minutes after irradiation, only dog 9 experienced decrement severe enough to be construed as incapacitation. Further, only dogs 5, 6, and 9 had severe decrement in two or more consecutive 10-trial sessions. At lower doses the severe decrement generally occurred during periods of vomiting; at higher doses, severe decrement also occurred during periods of convulsions.

As shown in Figure 2, the dogs performed quite well for several days (48-72 hours) after 5200- and 7800-rad doses. After doses of 10,200 rads or more, severe central nervous system damage (convulsions) was evident in most dogs, and a rapid decline in performance occurred within a few hours after irradiation.









Figure 2. Beagle performance after irradiation. Each vertical bar represents the average performance for each test session:  correct and  incorrect responses;  omissions. Each data point is an average of a 10-trial session. The standard deviation around the mean of the preirradiation base-line performance is indicated by the shaded area.




Figure 2 (continued)

Table II. Beagle Performance and Clinical Symptoms during the First Hour after Irradiation

Dog #	Midline tissue dose (rads)	Severe performance decrement (P.D.) ≤ 60% correct response			Number of 10-trial sessions with any P.D.*	Onset of vomiting (minutes)	Onset of convulsion (minutes)
		Time of severe P.D. (minutes)	Percent correct response during severe P.D.	Number of 10-trial sessions with severe P.D.			
1	5200	None	--	0	5	11	None
2	5200	30-32 45-47	40 50	2	7	10	None
3	7800	None	--	0	7	12	None
4	7800	10-12 33-35	30 60	2	5	10	None
5	10,200	12-15	35	2	3	7	None
6	10,200	0-4	55	2	10	30	60
7	11,900	0-2 54 →	30 60	3	8	None	~120
8	17,600	8-35 40-52	16 30	21	24	10	38
9	17,600	0-7 15-17 40 →	10 60 19	15	19	8	40

\* < 90% correct response for all dogs except dog 7 which was < 80%

The onset of complete incapacitation varied among dogs from a few hours to as long as 1 day before death.

#### IV. DISCUSSION

The pattern of trained beagle performance after irradiation is generally consistent with the results of Pitchford<sup>5</sup> and Block et al;<sup>1</sup> a definite early transient performance decrement was not consistently evident.



Although nearly all dogs experienced random periods of early performance decrement, these periods generally lasted only 1 to 3 minutes; they were not prolonged as was characteristic of the temporary debilitation observed in the trained monkeys, miniature pigs, and rats.

The decline in performance observed in a few of the dogs immediately after irradiation (Figure 2) may have been related to the noises involved in reactor operation. These dogs appeared distracted and usually responded incorrectly or omitted several consecutive trials. This transient performance decrement was not associated with the severe clinical symptoms (convulsions, coma, shock, etc.) that consistently occurred during the temporary performance decrement experienced by irradiated monkeys, pigs, and rats. When convulsions did occur in the dog, the animal could no longer perform and died shortly thereafter.

The dogs generally performed quite well until they were unable to stand in the testing box. At doses below 10,000 rads this occurred 3 to 4 days after irradiation and within 1 day of death. At higher doses, the early incidence of convulsions prevented satisfactory performance even though many of the dogs appeared to be alert and tried to perform while convulsing. The dogs performed reasonably well between the convulsive episodes; note the cyclic performance pattern shown by dog 8, Figure 2. Unlike monkeys, pigs, and rats, the dogs never appeared to lose consciousness until they became comatose a few hours before death.

Although three of the dogs were sacrificed, the survival time data that were obtained in this study agree reasonably well with survival time data obtained by Pitchford<sup>5</sup> and Block et al.<sup>1</sup>

## V. CONCLUSION

The response of trained dogs to supralethal doses of radiation is quite different from that of trained monkeys, miniature pigs, and rats. No sustained early temporary performance decrement was found; instead, very brief periods of performance decrement occurred randomly throughout the first 60 minutes after irradiation. At doses below 10,000 rads the animals continued to perform at or near satisfactory levels for several days after irradiation. Thereafter, a rapid decline in performance led to complete incapacitation within 1 day of death. When the dose exceeded 10,000 rads, central nervous system damage was obvious (convulsions), and the animals experienced severe decrement within 1 hour after irradiation; a few hours later, a rapid, irreversible decline in performance led to complete incapacitation and death.

## REFERENCES

1. Block, P. C., Taylor, J. F. and Ainsworth, E. J. Survival time and dose-response following supralethal irradiation of dogs. *Radiation Res.* 45:384-398, 1971.
2. Casarett, A. P. and Comar, C. L. Incapacitation and performance decrement following split doses of fission spectrum radiation. Ithaca, New York, Department of Physical Biology, New York State Veterinary College, Cornell University (Final Summary Report, Contract No. DASA-01-68-C-0162), 1970.
3. Chaput, R. L. and Wise, D. Miniature pig incapacitation and performance decrement after mixed gamma-neutron irradiation. *Aerospace Medicine* 41:290-293, 1970.
4. Curran, C. R., Conrad, D. W. and Young, R. W. The effects of 2000 rads of pulsed gamma-neutron radiation upon the performance of unfettered monkeys. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR71-3, 1971.
5. Pitchford, T. L. Beagle incapacitation and survival time after pulsed mixed gamma-neutron irradiation. Bethesda, Maryland, Armed Forces Radiobiology Research Institute Scientific Report SR68-24, 1968.

## DISTRIBUTION LIST

### AIR FORCE

The Surgeon General, U.S. Department of the Air Force, Washington, D. C. 20314 (1)  
Executive Officer, Director of Professional Services, Office of the Surgeon General, Hq. USAF (AFMSPA),  
Washington, D. C. 20314 (1)  
Headquarters, U. S. Air Force (AFMSPAB), Washington, D. C. 20314 (1)  
Office of the Command Surgeon (ADCSG), Hq. ADC, USAF, Ent AFB, Colorado 80912 (1)  
Air Force Weapons Laboratory, ATTN: WLIL (1), ATTN: WLRB-2 (1), Kirtland AFB, New Mexico 87117 (2)  
Chief, Nuclear Medicine Department, P. O. Box 5088, USAF Hospital, Wright-Patterson AFB, Ohio 45433 (1)  
USAFSAM (RA), ATTN: Chief, Radiobiology Division, Brooks AFB, Texas 78235 (1)  
USAFSAM, ATTN: EDAD, Brooks AFB, Texas 78235 (1)

### ARMY

The Surgeon General, U. S. Department of the Army, Washington, D. C. 20314 (1)  
Surgeon General, ATTN: MEDDH-N, U. S. Department of the Army, Washington, D. C. 20314 (1)  
USACDC CSSG, Doctrine Division, Fort Lee, Virginia 23801 (1)  
Commanding Officer, USACDC CBR Agency, Fort McClellan, Alabama 36201 (1)  
Commanding Officer, U. S. Army Combat Developments Command, Institute of Nuclear Studies, Fort Bliss, Texas  
79916 (1)  
CG, USCONARC, ATTN: ATUTR-TNG (NBC), Fort Monroe, Virginia 23351 (1)  
Nuclear Branch AMCRD-DN-RE, U. S. Army Materiel Command, Washington, D. C. 20315 (1)  
Commanding Officer, U. S. Army Medical Research Laboratory, Fort Knox, Kentucky 40121 (1)  
Commanding Officer, USA Nuclear Medical Research Detachment, Europe, APO New York, N. Y. 09180 (2)  
Chief of Research and Development, ATTN: Nuclear, Chemical and Biological Division, U. S. Department of the  
Army, Washington, D. C. 20310 (1)  
Army Research Office, ATTN: Chief, Scientific Analysis Branch, Life Sciences Division, 3045 Columbia Pike,  
Arlington, Virginia 22204 (1)  
Division of Nuclear Medicine, Walter Reed Army Institute of Research, Walter Reed Army Medical Center,  
Washington, D. C. 20012 (5)  
Commanding Officer, U. S. Army Environmental Hygiene Agency, ATTN: USAEHA-RP, Edgewood Arsenal, Mary-  
land 21010 (1)  
Commandant, U. S. Army Medical Field Service School, ATTN: MEDEW-ZNW, Fort Sam Houston, Texas 78234 (1)

### NAVY

Chief, Bureau of Medicine and Surgery, U. S. Navy Department, Washington, D. C. 20390 (1)  
Chief, Bureau of Medicine and Surgery, ATTN: Code 71, U. S. Navy Department, Washington, D. C. 20390 (1)  
Commanding Officer, Naval Aerospace Medical Institute, Naval Aviation Medical Center, ATTN: Director of  
Research, Pensacola, Florida 32512 (3)  
Commanding Officer, Nuclear Weapons Training Center, Atlantic, Nuclear Warfare Department, Norfolk, Virginia  
23511 (1)  
Commanding Officer, Nuclear Weapons Training Center, Pacific, U. S. Naval Air Station, North Island, San Diego,  
California 92135 (1)  
Director, Biological Sciences Division, Office of Naval Research, Washington, D. C. 20360 (1)  
Commanding Officer, U. S. Naval Hospital, ATTN: Director, REEL, National Naval Medical Center, Bethesda,  
Maryland 20014 (1)  
Head, Animal Behavioral Sciences Branch, Naval Aerospace Medical Institute, Naval Aerospace Medical Center,  
ATTN: Dr. John S. Thach, Jr., Pensacola, Florida 32512 (1)  
Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base, NL, ATTN: Medical Library,  
Groton, Connecticut 06340 (1)  
Commanding Officer, Naval Submarine Medical Center, Naval Submarine Base, NL, ATTN: Code 53, Nuclear Medi-  
cine Training Division, Groton, Connecticut 06340 (1)

### D. O. D.

Director, Defense Nuclear Agency, Washington, D. C. 20305 (1)  
Director, Defense Nuclear Agency, ATTN: DDST, Washington, D. C. 20305 (1)  
Director, Defense Nuclear Agency, ATTN: Mr. P. H. Haas, DDST, Washington, D. C. 20305 (1)  
Director, Defense Nuclear Agency, ATTN: Chief, Radiation Directorate, Washington, D. C. 20305 (1)

D. O. D. (continued)

Director, Defense Nuclear Agency, ATTN: Technical Library (APTL), Washington, D. C. 20305 (2)  
Commanding Officer, Harry Diamond Laboratories, ATTN: Nuclear Vulnerability Branch, Washington, D. C. 20438 (1)  
Commander, Field Command, Defense Nuclear Agency, ATTN: FC Technical Library, Kirtland AFB, New Mexico 87117 (1)  
Commander, Headquarters Field Command, Defense Nuclear Agency, ATTN: FCTG8, Kirtland AFB, New Mexico 87117 (2)  
Director, Armed Forces Institute of Pathology, Washington, D. C. 20305 (1)  
Administrator, Defense Documentation Center, Cameron Station, Bldg. 5, Alexandria, Virginia 22314 (12)

OTHER GOVERNMENT

U. S. Atomic Energy Commission, Headquarters Library, Reports Section, Mail Station G-17, Washington, D. C. 20545 (1)  
U. S. Atomic Energy Commission, Division of Biology and Medicine, Washington, D. C. 20545 (1)  
U. S. Atomic Energy Commission, Bethesda Technical Library, 7920 Norfolk Avenue, Bethesda, Maryland 20014 (1)  
National Aeronautics and Space Administration, Manned Spacecraft Center, ATTN: Dr. Charles M. Barnes, Code DC7, Houston, Texas 77058 (1)  
National Aeronautics and Space Administration, Manned Spacecraft Center, ATTN: Dr. B. D. Newsom, Code DB4, Houston, Texas 77058 (1)  
National Bureau of Standards, ATTN: Chief, Radiation Physics Division, Washington, D. C. 20234 (1)  
U. S. Public Health Service, Bureau of Radiological Health, Division of Biological Effects, 12720 Twinbrook Parkway, Rockville, Maryland 20852 (1)  
U. S. Public Health Service, Bureau of Radiological Health, Library, 12720 Twinbrook Parkway, Rockville, Maryland 20852 (1)  
U. S. Public Health Service, Northeastern Radiological Health Laboratory, 109 Holton Street, Winchester, Massachusetts 01890 (1)  
U. S. Public Health Service, Southeastern Radiological Health Laboratory, P. O. Box 61, Montgomery, Alabama 36101 (1)  
U. S. Public Health Service, Southwestern Radiological Health Laboratory, P. O. Box 15027, Las Vegas, Nevada 89114 (1)

OTHER

Argonne National Laboratory, Library Services Department, Report Section Bldg. 203, RM-CE-125, 9700 South Cass Avenue, Argonne, Illinois 60440 (1)  
Dr. Donald G. Baker, Radiobiology Department, Zellerbach Saroni Tumor Institute, 1600 Divisadero Street, San Francisco, California 94115 (1)  
Dr. J. T. Brennan, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pennsylvania 19104 (1)  
Brookhaven National Laboratory, Information Division, ATTN: Research Library, Upton, Long Island, New York 11973 (2)  
Dr. J. S. Burkle, Director of Nuclear Medicine, York Hospital, York, Pennsylvania 17403 (1)  
S. C. Bushong, Department of Radiology, Baylor University College of Medicine, Houston, Texas 77024 (1)  
University of California, Lawrence Radiation Laboratory, Library, Bldg. 50, Room 134, Berkeley, California 94720 (1)  
Director, Radiobiology Laboratory, University of California, Davis, California 95616 (1)  
University of California, Lawrence Radiation Laboratory, Technical Information Division Library L-3, P. O. Box 808, Livermore, California 94551 (2)  
University of California, Laboratory of Nuclear Medicine and Radiation Biology, Library, 900 Veteran Avenue, Los Angeles, California 90024 (1)  
Dr. C. Jelleff Carr, Director, Life Sciences Research Office, Federation of American Societies for Experimental Biology, 9650 Rockville Pike, Bethesda, Maryland 20014 (1)  
Cdr. William H. Chapman, USN (Ret.), Bio-Medical Division L-523, Lawrence Radiation Laboratory, University of California, P. O. Box 808, Livermore, California 94551 (1)  
Director, Collaborative Radiological Health Laboratory, Colorado State University, Fort Collins, Colorado 80521 (1)  
Dr. L. W. Davis, Radiology Department, University of Pennsylvania, 3400 Spruce Street, Philadelphia, Pennsylvania 19104 (1)  
Professor Merril Eisenbud, New York University, Tuxedo, New York 10987 (1)  
Dr. T. C. Evans, Radiation Research Laboratory, College of Medicine, University of Iowa, Iowa City, Iowa 52240 (1)



#### OTHER (continued)

- Dr. Arnold Feldman, Institute of Radiology, School of Medicine, Washington University, 510 South Kingshighway, St. Louis, Missouri 63110 (1)
- Mr. Orin Gelderloos, Division of Literature, University of Michigan, Dearborn Campus, Dearborn, Michigan 48124 (1)
- General Dynamics/Fort Worth, ATTN: Librarian, P. O. Box 748, Fort Worth, Texas 76101 (1)
- Gulf General Atomic Incorporated, ATTN: Library, P. O. Box 608, San Diego, California 92112 (1)
- Dr. James E. Huff, Department of Pharmacology and Toxicology, School of Medicine and Dentistry, University of Rochester, 260 Crittenden Blvd., Rochester, New York 14620 (1)
- IIT Research Institute, ATTN: Document Library, 10 West 35th Street, Chicago, Illinois 60616 (1)
- Johns Hopkins University, Applied Physics Laboratory, ATTN: Document Library, 8621 Georgia Avenue, Silver Spring, Maryland 20910 (1)
- Dr. R. F. Kallman, Department of Radiology, Stanford University, Palo Alto, California 94305 (1)
- Dr. L. S. Kelly, Donner Laboratory, University of California at Berkeley, Berkeley, California 94720 (1)
- Dr. Robert Landolt, Bionucleonics Department, Purdue University, Lafayette, Indiana 47907 (1)
- Los Alamos Scientific Laboratory, ATTN: Report Librarian, P. O. Box 1663, Los Alamos, New Mexico 87544 (1)
- Director, Nuclear Science Center, Louisiana State University, Baton Rouge, Louisiana 70803 (2)
- Lovelace Foundation for Medical Education and Research, Document Library, 5200 Gibson Blvd., S. E., Albuquerque, New Mexico 87108 (1)
- Dr. Ross A. McFarland, Guggenheim Professor of Aerospace Health and Safety, Harvard School of Public Health, 665 Huntington Avenue, Boston, Massachusetts 02115 (1)
- Dr. J. I. Marcum, Rand Corporation, 1700 Main Street, Santa Monica, California 90401 (1)
- Massachusetts Institute of Technology, M. I. T. Libraries, Technical Reports, Room 14 E-210, Cambridge, Massachusetts 02139 (1)
- Dr. Charles W. Mays, Physics Group Leader, Radiobiology Division, University of Utah, Salt Lake City, Utah 84112 (1)
- Ohio State University, Nuclear Reactor Laboratory, 1298 Kinnear Road, Columbus, Ohio 43212 (1)
- Dr. Harvey M. Patt, Laboratory of Radiobiology, University of California, San Francisco Medical Center, San Francisco, California 94122 (1)
- Purdue University, Nuclear Engineering Library, Lafayette, Indiana 47907 (1)
- Dr. S. M. Reichard, Director, Division of Radiobiology, Medical College of Georgia, Augusta, Georgia 30902 (1)
- Dr. H. H. Rossi, 630 West 168th Street, New York, N. Y. 10032 (1)
- Dr. Eugene L. Saenger, Director, Radioisotope Laboratory, Cincinnati General Hospital, Cincinnati, Ohio 45229 (1)
- Sandia Corporation Library, P. O. Box 5800, Albuquerque, New Mexico 87115 (1)
- Scientific Committee on the Effects of Atomic Radiation, ATTN: Library, United Nations Room 3267, United Nations Plaza, New York, N. Y. 10017 (1)
- Scope Publications, Franklin Station, P. O. Box 7407, Washington, D. C. 20004 (1)
- Texas A and M University, Radiation Biology Laboratory, Texas Engineering Experiment Station, College Station, Texas 77840 (2)
- Texas Nuclear Corporation, ATTN: Director of Research, Box 9267 Allandale Station, Austin, Texas 78756 (1)
- University of Rochester, Atomic Energy Project Library, P. O. Box 287, Station 3, Rochester, New York 14620 (1)
- University of Southern California, Nuclear Physics Laboratory, University Park, Los Angeles, California 90007 (1)
- Western Reserve University, Department of Radiology, Division of Radiation Biology, Cleveland, Ohio 44106 (1)
- Mr. Lionel Zamore, 601 Brightwater Court, Brooklyn, New York 11235 (1)

#### FOREIGN

- International Atomic Energy Agency, Kärrntnerring 11, Vienna I. 1010, Austria (1)
- European Atomic Energy Community, C. E. E. A., Library, 51 rue Belliard, Brussels 4, Belgium (1)
- Dr. L. G. Lajtha, Paterson Laboratories, Christie Hospital and Holt Radium Institute, Manchester, England (1)
- Dr. L. F. Lamerton, Biophysics Department, Institute of Cancer Research, Surrey Branch, Belmont, Sutton, Surrey, England (1)
- National Lending Library for Science and Technology, Boston Spa, Yorkshire, England (1)
- Directorate of Medical and Health Services, FAF (Federal Armed Forces), Bonn, Ermekeilstrasse 27, West Germany (1)
- Abteilung für Strahlenbiologie im Institut für Biophysik der Universität Bonn, 53 Bonn-Venusberg, Annaberger Weg 15, Federal Republic of Germany (2)
- Prof. Dr. H. Langendorff, Direktor des Radiologischen Instituts der Universität, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)

FOREIGN (continued)

- Priv.-Doz. Dr. O. Messerschmidt, Radiologisches Institut der Universität, 78 Freiburg im Breisgau, Albertstrasse 23, Germany (1)
- Dr. Helmut Mitschrich, Sanitätsamt der Bundeswehr, 53 Bonn-Beuel, Zingsheimstrasse 5, Germany (2)
- Prof. Dr. F. Wachsmann, Gesellschaft für Strahlenforschung m.b.H., 8042 Neuherberg bei München, Institut für Strahlenschutz, Ingolstadter Landstrasse 1, München, Germany (1)
- Dr. M. Feldman, Section of Cell Biology, The Weizmann Institute of Science, Rehovoth, Israel (1)
- Dr. G. W. Barendsen, Radiobiological Institute TNO, Rijswijk, Netherlands (1)
- Dr. L. M. van Putten, Radiobiological Institute TNO, 151 Lance Kleiweg, Rijswijk 2 H, Netherlands (1)
- Puerto Rico Nuclear Center, ATTN: Reading Room, College Station, Mayaguez, Puerto Rico 00708 (2)
- Dr. H. Cottier, Pathological Institut der Universität, Bern, Switzerland (1)

UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - R &amp; D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Armed Forces Radiobiology Research Institute Defense Nuclear Agency Bethesda, Maryland 20014		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
		2b. GROUP N/A	
3. REPORT TITLE  PERFORMANCE OF TRAINED BEAGLES AFTER SUPRALETHAL DOSES OF RADIATION			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name)  R. L. Chaput, R. T. Kovacic and E. L. Barron			
6. REPORT DATE February 1972		7a. TOTAL NO. OF PAGES 25	7b. NO. OF REFS 5
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)  AFRRI SR72-1	
b. PROJECT NO. NWER XAXM			
c. Task and Subtask A 904		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d. Work Unit 05			
10. DISTRIBUTION STATEMENT  Approved for public release; distribution unlimited			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY Director Defense Nuclear Agency Washington, D. C. 20305	
13. ABSTRACT  Nine beagle dogs were trained by shock avoidance conditioning to work a discrimination problem cued by visual and auditory stimuli. The animals received 5200- to 17,600-rad midline tissue doses of pulsed mixed gamma-neutron radiation, and their postirradiation performance was evaluated. During the first 60 minutes after irradiation, random, brief periods of performance decrement frequently occurred. Dogs that received less than 10,000 rads performed at or near preirradiation levels several days until they became permanently incapacitated shortly before death. After doses of 10,000 rads or more most dogs performed satisfactorily for only a few minutes to hours. Their condition then worsened precipitously and the dogs died shortly thereafter.			